

BIOLOGY *of* REPRODUCTION

Official Journal of the Society for
the Study of Reproduction

JANUARY 2007

VOLUME 76 NUMBER 1

| | |
|---|----|
| Biology of Reproduction Highlights | 1 |
| The Expression of Multiple Connexins Throughout Spermatogenesis in the Rainbow Trout Testis Suggests a Role for Complex Intercellular Communication | 2 |
| <i>Benjamin de Montgolfier, Julie Dufresne, Myriam Letourneau, James J. Nagler, Alain Fournier, Céline Audet, and Daniel G. Cyr</i> | |
| Intercellular communication in the fish testis is complex, involving several different connexins, and is a highly regulated process. | |
| Heterocomplex Formation and Cell-Surface Accumulation of Hen's Serum Zona Pellucida B1 (ZPB1) with ZPC Expressed by a Mammalian Cell Line (COS-7): A Possible Initiating Step of Egg-Envelope Matrix Construction | 9 |
| <i>Hiroki Okumura, Naohito Aoki, Chihiro Sato, Daita Nadano, and Tsukasa Matsuda</i> | |
| Hen's serum ZPB1 specifically bound to ZPC expressed and secreted by COS-7 cells through its C-terminal region of ZP-domain and accumulated as ZPB1-ZPC complexes on the surface of the COS-7 cells. | |
| Regulation of Cavernous Nerve Injury-Induced Apoptosis by Sonic Hedgehog | 19 |
| <i>Carol A. Podlasek, Cynthia L. Meroz, Yi Tang, Kevin E. McKenna, and Kevin T. McVary</i> | |
| SHH treatment at the time of CN injury prevented CN injury-induced apoptosis in a dose-dependent manner, suggesting that SHH stabilizes alterations of the corpus cavernosal smooth muscle following CN injury. | |
| Establishment of a Functional Ovine Fetoplacental Artery Endothelial Cell Line with a Prolonged Life Span | 29 |
| <i>Yang Song and Jing Zheng</i> | |
| Establishment of a functional ovine fetoplacental artery endothelial cell line with an extended life span provides a valuable cell model for studying fetoplacental endothelial function. | |
| Epigenetic Marks in Cloned Rhesus Monkey Embryos: Comparison with Counterparts Produced In Vitro | 36 |
| <i>Jifeng Yang, Shihua Yang, Nathalie Beaujean, Yuyu Niu, Xiechao He, Yunhua Xie, Xianghui Tang, Liu Wang, Qi Zhou, and Weizhi Ji</i> | |
| Global epigenetic reprogramming, such as DNA methylation and histone acetylation, was always abnormal in cloned rhesus embryos. | |
| Building a Testis: Formation of Functional Testis Tissue after Transplantation of Isolated Porcine (<i>Sus scrofa</i>) Testis Cells | 43 |
| <i>Ali Honaramooz, Susan O. Megee, Rahul Rathi, and Ina Dobrinski</i> | |
| Isolated cells from neonatal porcine testis form functional testis tissue with complete spermatogenesis after ectopic xenotransplantation into mouse hosts assisted reproductive technology, germ cells, morphogenesis, pig, spermatogenesis, testis. | |
| In the Absence of the Mouse DNA/RNA-Binding Protein MSY2, Messenger RNA Instability Leads to Spermatogenic Arrest | 48 |
| <i>Juxiang Yang, Carlos R. Morales, Sergey Medvedev, Richard M. Schultz, and Norman B. Hecht</i> | |
| In the absence of MSY2, transcription, mRNA processing, and transport are normal, but a precocious redistribution of mRNA onto polysomes and decreases in many germ cell mRNAs indicate RSY2 is essential for mRNA storage and stabilization. | |
| Leukemia Inhibitory Factor Enhances Formation of Germ Cell Colonies in Neonatal Mouse Testis Culture | 55 |
| <i>Mito Kanatsu-Shinohara, Kimiko Inoue, Narumi Ogonuki, Hiromi Miki, Shosei Yoshida, Shinya Toyokuni, Jiyoung Lee, Atsuo Ogura, and Takashi Shinohara</i> | |
| Leukemia inhibitory factor enhances germ cell colony formation. | |
| RNase9, an Androgen-Dependent Member of the RNase A Family, Is Specifically Expressed in the Rat Epididymis | 63 |
| <i>Chun-Fang Zhu, Qiang Liu, Li Zhang, Hai-Xin Yuan, Wei Zhen, Jin-Song Zhang, Zheng-Jun Chen, Susan H. Hall, Frank S. French, and Yong-Lian Zhang</i> | |
| Rat RNase9, an androgen-dependent member of the RNase A superfamily, is exclusively expressed in epididymis, abundantly secreted from the caput principal cells into lumen and bound on the sperm acrosome, suggesting roles in sperm maturation. | |
| Production of Recombinant Orange-Spotted Grouper (<i>Epinephelus coioides</i>) Luteinizing Hormone in Insect Cells by the Baculovirus Expression System and Its Biological Effect | 74 |
| <i>Miao Cui, Wensheng Li, Wen Liu, Kai Yang, Yi Pang, and Lin Haoran</i> | |
| Recombinant grouper Lh was expressed by the baculovirus-insect system, and its biological activity was in accord with the long positive feedback loop of Lh on sex steroid hormones and the short negative feedback on gnRH mRNA levels. | |

| | |
|--|-----|
| The Proximal <i>Gata4</i> Promoter Directs Reporter Gene Expression to Sertoli Cells During Mouse Gonadal Development | 85 |
| <i>Séverine Mazaud Guittot, Amélie Tétu, Eric Legault, Nicolas Pilon, David W. Silversides, and Robert S. Viger</i> | |
| The first 5 kb of 5' flanking region of the <i>Gata4</i> gene is sufficient to drive transgene expression in the testis, and specifically Sertoli cells, from the onset of gonadal morphogenesis to adulthood. | |
| Functional HY-Specific CD8⁺ T Cells Are Found in a High Proportion of Women Following Pregnancy with a Male Fetus | 96 |
| <i>Karen P. Piper, Andrew McLarnon, Julie Arrazi, Claire Horlock, Jennifer Ainsworth, Mark D. Kilby, William L. Martin, and Paul A. Moss</i> | |
| Alloreactive CD8 ⁺ T cells are generated frequently following normal pregnancy and retain functional capability for years following pregnancy. | |
| Decidual Stromal Cell Response to Paracrine Signals from the Trophoblast: Amplification of Immune and Angiogenic Modulators | 102 |
| <i>A.P. Hess, A.E. Hamilton, S. Talbi, C. Dosiou, M. Nyegaard, N. Nayak, O. Genbecev-Krtolica, P. Mavrogianis, K. Ferrer, J. Kruessel, A.T. Fazleabas, S.J. Fisher, and L.C. Giudice</i> | |
| Decidualized human endometrial fibroblasts respond to trophoblast-secreted products by acquisition of an immune and proangiogenic phenotype. | |
| Vascular Hyperresponsiveness to Adrenomedullin During Pregnancy Is Associated with Increased Generation of Cyclic Nucleotides in Rat Mesenteric Artery | 118 |
| <i>Gracious R. Ross and Chandra Yallampalli</i> | |
| Vasodilatory effects of adrenomedullin are greater during pregnancy and are associated with increased generation of cyclic nucleotides in resistance vessels, and this may contribute to cardiovascular adaptations. | |
| The Expression of Prostaglandin Endoperoxide Synthase 2 Messenger RNA and the Proportion of Smooth Muscle and Collagen in the Sheep Cervix During the Estrous Cycle | 124 |
| <i>Claire M. Kershaw, Rex J. Scaramuzzi, Michael R. McGowan, Caroline P.D. Wheeler-Jones, and Muhammad Khalid</i> | |
| Prostaglandin endoperoxide synthase 2 mRNA expression is greater in the sheep cervix at estrus compared to other stages of the estrous cycle and may regulate cervical relaxation through the separation of collagen fibers and bundles. | |
| CDH1 Is a Specific Marker for Undifferentiated Spermatogonia in Mouse Testes | 130 |
| <i>Masutaka Tokuda, Yuzo Kadokawa, Hiroki Kurahashi, and Tohru Marunouchi</i> | |
| CDH1 is specifically expressed in undifferentiated type A spermatogonia, which contain spermatogenic stem cells, in mouse testes. | |
| 17Beta-Estradiol Triggers Postspawning in Spermatogenically Active Gilthead Seabream (<i>Sparus aurata</i> L.) Males | 142 |
| <i>E. Chaves-Pozo, S. Liarte, L. Vargas-Chacoff, A. García-López, V. Mulero, J. Mesequer, J.M. Mancera, and A. García-Ayala</i> | |
| Treating spermatogenically active gilthead seabream males with high doses of estradiol for short periods of time induced several of the processes involved in postspawning, including testicular regression and acidophilic granulocyte infiltration. | |
| Differential Expression of Genes Encoding Constitutive and Inducible 20S Proteasomal Core Subunits in the Testis and Epididymis of Theophylline- or 1,3-Dinitrobenzene-Exposed Rats | 149 |
| <i>Mark W. Tengowski, Dongyan Feng, Miriam Sutovsky, and Peter Sutovsky</i> | |
| Theophylline or 1,3-dinitrobenzene exposure produces similar morphologic and gene expression changes that can be measured in the ubiquitin-proteasome system as a possible marker for reproductive toxicology. | |
| The Cytotrophoblast Layer of Human Chorionic Villi Becomes Thinner but Maintains Its Structural Integrity During Gestation | 164 |
| <i>Miki Mori, Gen Ishikawa, Shan-Shun Luo, Takuya Mishima, Tadashi Goto, John M. Robinson, Shigeki Matsubara, Toshiyuki Takeshita, Hiroaki Kataoka, and Toshihiro Takizawa</i> | |
| Cytochemistry of hepatocyte growth factor activator inhibitor type 1, a new specific marker of the cytotrophoblast (CTB), reveals that the CTB layer of human chorionic villi becomes thinner but maintains its structural integrity during gestation. | |
| Identification, Culture, and Characterization of Germline Stem Cell-Like Cells in Chicken Testes | 173 |
| <i>Jin Gyoung Jung, Young Mok Lee, Tae Sub Park, Sang Hyun Park, Jeong Mook Lim, and Jae Yong Han</i> | |
| Germline stem cell-like cells were present in the chicken testicular cells, and these cells were positively stained with germ cell-specific markers. | |

Contents by Category

Embryo

- 36 Epigenetic Marks in Cloned Rhesus Monkey Embryos: Comparison with Counterparts Produced In Vitro. *Jifeng Yang, Shihua Yang, Nathalie Beaujean, Yuyu Niu, Xiechao He, Yunhua Xie, Xianghui Tang, Liu Wang, Qi Zhou, and Weizhi Ji*

Female Reproductive Tract

- 102 Decidual Stromal Cell Response to Paracrine Signals from the Trophoblast: Amplification of Immune and Angiogenic Modulators. *A.P. Hess, A.E. Hamilton, S. Talbi, C. Dosiou, M. Nyegaard, N. Nayak, O. Genbecev-Krtolica, P. Mavrogianis, K. Ferrer, J. Kruessel, A.T. Fazleabas, S.J. Fisher, and L.C. Giudice*
- 124 The Expression of Prostaglandin Endoperoxide Synthase 2 Messenger RNA and the Proportion of Smooth Muscle and Collagen in the Sheep Cervix During the Estrous Cycle. *Claire M. Kershaw, Rex J. Scaramuzzi, Michael R. McGowan, Caroline P.D. Wheeler-Jones, and Muhammad Khalid*

Gamete Biology

- 9 Heterocomplex Formation and Cell-Surface Accumulation of Hen's Serum Zona Pellucida B1 (ZPB1) with ZPC Expressed by a Mammalian Cell Line (COS-7): A Possible Initiating Step of Egg-Envelope Matrix Construction. *Hiroki Okumura, Naohito Aoki, Chihiro Sato, Daita Nadano, and Tsukasa Matsuda*

Immunology

- 96 Functional HY-Specific CD8⁺ T Cells Are Found in a High Proportion of Women Following Pregnancy with a Male Fetus. *Karen P. Piper, Andrew McLarnon, Julie Arrazi, Claire Horlock, Jennifer Ainsworth, Mark D. Kilby, William L. Martin, and Paul A. Moss*

Male Reproductive Tract

- 19 Regulation of Cavemous Nerve Injury-Induced Apoptosis by Sonic Hedgehog. *Carol A. Podlasek, Cynthia L. Meroz, Yi Tang, Kevin E. McKenna, and Kevin T. McVary*
- 63 RNase9, an Androgen-Dependent Member of the RNase A Family, Is Specifically Expressed in the Rat Epididymis. *Chun-Fang Zhu, Qiang Liu, Li Zhang, Hai-Xin Yuan, Wei Zhen, Jin-Song Zhang, Zheng-Jun Chen, Susan H. Hall, Frank S. French, and Yong-Lian Zhang*

Mechanisms of Hormone Action

- 74 Production of Recombinant Orange-Spotted Grouper (*Epinephelus coioides*) Luteinizing Hormone in Insect Cells by the Baculovirus Expression System and Its Biological Effect. *Miao Cui, Wensheng Li, Wen Liu, Kai Yang, Yi Pang, and Lin Haoran*

Pregnancy

- 29 Establishment of a Functional Ovine Fetoplacental Artery Endothelial Cell Line with a Prolonged Life Span. *Yang Song and Jing Zheng*

- 118 Vascular Hyperresponsiveness to Adrenomedullin During Pregnancy Is Associated with Increased Generation of Cyclic Nucleotides in Rat Mesenteric Artery. *Gracious R. Ross and Chandra Yallampalli*
- 164 The Cytotrophoblast Layer of Human Chorionic Villi Becomes Thinner but Maintains Its Structural Integrity During Gestation. *Miki Mori, Gen Ishikawa, Shan-Shun Luo, Takuya Mishima, Tadashi Goto, John M. Robinson, Shigeki Matsubara, Toshiyuki Takeshita, Hiroaki Kataoka, and Toshihiro Takizawa*

Reproductive Technology

- 43 Building a Testis: Formation of Functional Testis Tissue after Transplantation of Isolated Porcine (*Sus scrofa*) Testis Cells. *Ali Honaramooz, Susan O. Megee, Rahul Rath, and Ina Dobrinski*
- 55 Leukemia Inhibitory Factor Enhances Formation of Germ Cell Colonies in Neonatal Mouse Testis Culture. *Mito Kanatsu-Shinohara, Kimiko Inoue, Narumi Ogonuki, Hiromi Miki, Shosei Yoshida, Shinya Toyokuni, Jiyoung Lee, Atsuo Ogura, and Takashi Shinohara*

Testis

- 2 The Expression of Multiple Connexins Throughout Spermatogenesis in the Rainbow Trout Testis Suggests a Role for Complex Intercellular Communication. *Benjamin de Montgolfier, Julie Dufresne, Myriam Letourneau, James J. Nagler, Alain Fournier, Céline Audet, and Daniel G. Cyr*
- 48 In the Absence of the Mouse DNA/RNA-Binding Protein MSY2, Messenger RNA Instability Leads to Spermatogenic Arrest. *Juxiang Yang, Carlos R. Morales, Sergey Medvedev, Richard M. Schultz, and Norman B. Hecht*
- 85 The Proximal *Gata4* Promoter Directs Reporter Gene Expression to Sertoli Cells During Mouse Gonadal Development. *Séverine Mazaud Guittot, Amélie Tétu, Eric Legault, Nicolas Pilon, David W. Silversides, and Robert S. Viger*
- 130 CDH1 Is a Specific Marker for Undifferentiated Spermatogonia in Mouse Testes. *Masutaka Tokuda, Yuzo Kadokawa, Hiroki Kurahashi, and Tohru Marunouchi*
- 142 17Beta-Estradiol Triggers Postspawning in Spermatogenically Active Gilthead Seabream (*Sparus aurata* L.) Males. *E. Chaves-Pozo, S. Liarte, L. Vargas-Chacoff, A. García-López, V. Mulero, J. Meseguer, J.M. Mancera, and A. García-Ayala*
- 173 Identification, Culture, and Characterization of Germline Stem Cell-Like Cells in Chicken Testes. *Jin Gyoung Jung, Young Mok Lee, Tae Sub Park, Sang Hyun Park, Jeong Mook Lim, and Jae Yong Han*

Toxicology

- 149 Differential Expression of Genes Encoding Constitutive and Inducible 20S Proteasomal Core Subunits in the Testis and Epididymis of Theophylline- or 1,3-Dinitrobenzene-Exposed Rats. *Mark W. Tengowski, Dongyan Feng, Miriam Sutovsky, and Peter Sutovsky*