

BIOLOGY *of* REPRODUCTION

Official Journal of the Society for
the Study of Reproduction

JULY 2007

VOLUME 77 NUMBER 1

Biology of Reproduction Highlights	1
Oxidative Metabolism of Pyruvate Is Required for Meiotic Maturation of Murine Oocytes In Vivo	2
<i>Mark T. Johnson, Edward A. Freeman, David K. Gardner, and Patricia A. Hunt</i> Ablation of the <i>Pdhal</i> subunit gene of pyruvate dehydrogenase in developing oocytes compromises meiotic maturation but not growth or ovulation.	
Impact of Dietary Fatty Acids on Oocyte Quality and Development in Lactating Dairy Cows	9
<i>Ali A. Fouladi-Nashta, Carlos G. Gutierrez, Jin G. Gong, Philip C. Garnsworthy, and Robert Webb</i> Increased level of dietary fatty acids improved developmental potential of oocytes to the blastocyst stage and also improved blastocyst quality.	
Insulin-Like Growth Factor (IGF) 2 Stimulates Steroidogenesis and Mitosis of Bovine Granulosa Cells Through the IGF1 Receptor: Role of Follicle-Stimulating Hormone and IGF2 Receptor	18
<i>L.J. Spicer and P.Y. Aad</i> IGF2 increases proliferation and stimulates differentiation in granulosa cells via the IGF1 receptor, and the IGF2 receptor may be hormonally regulated and associated with decreased granulosa cell responses.	
Fibroblast Growth Factor 2 Is More Dynamic than Vascular Endothelial Growth Factor A During the Follicle-Luteal Transition in the Cow	28
<i>R.S. Robinson, L.T. Nicklin, A.J. Hammond, D. Schams, M.G. Hunter, and G.E. Mann</i> Follicular FGF2 is up-regulated during the preovulatory period by the LH surge and plays an active role in initiation of luteal angiogenesis.	
Catsper3 and Catsper4 Are Essential for Sperm Hyperactivated Motility and Male Fertility in the Mouse	37
<i>Jingling Jin, Nange Jin, Huili Zheng, Seungil Ro, Dora Tafolla, Kenton M. Sanders, and Wei Yan</i> <i>Catsper3</i> and <i>Catsper4</i> are essential for sperm hyperactivated motility and male fertility.	
Modulation of Human Arterial Tone During Pregnancy: The Effect of the Bioactive Metabolite Sphingosine-1-Phosphate	45
<i>Nicola K. Hudson, Maureen O'Hara, Helen A. Lacey, Jemma Corcoran, Denise G. Hemmings, Mark Wareing, Philip Baker, and Michael J. Taggart</i> Sphingosine-1-phosphate produces dose-dependent vasoconstriction in human omental and myometrial arteries during pregnancy, with tissue specific mechanisms of action.	
Epidermal Growth Factor-Like Growth Factors Prevent Apoptosis of Alcohol-Exposed Human Placental Cytotrophoblast Cells	53
<i>Garen S. Wolff, Po Jen Chiang, Susan M. Smith, Roberto Romero, and D. Randall Armant</i> Physiologically relevant levels of ethanol rapidly induce human cytotrophoblast cells to apoptosis; however, endogenous production of the survival factor HBEGF could provide the placenta a protective mechanism during prenatal alcohol exposure.	
Actions of Epidermal Growth Factor Receptor/Mitogen-Activated Protein Kinase and Protein Kinase C Signaling in Granulosa Cells from <i>Gallus gallus</i> Are Dependent upon Stage of Differentiation	61
<i>Dori C. Woods, Morgan J. Haugen, and A.L. Johnson</i> The actions of EGFR/MAPK and protein kinase C signaling on granulosa cell steroidogenesis are dependent upon stage of differentiation.	
Factors Maintaining Normal Sperm Tail Structure During Epididymal Maturation Studied in <i>Gopc</i>^{-/-} Mice	71
<i>Fumie Suzuki-Toyota, Chizuru Ito, Yoshiro Toyama, Mamiko Maekawa, Ryoji Yao, Tetsuo Noda, Hiroshi Iida, and Kiyotaka Toshimori</i> Tails disorganize during epididymal maturation of <i>Gopc</i> ^{-/-} spermatozoa, suggesting the posterior ring, the connecting piece, SPATA19/spergen-1, and GPX4/PHGPx are requisite for maintaining the normal tail morphology in the epididymis.	
Signaling Pathways for Germ Cell Death in Adult Cynomolgus Monkeys (<i>Macaca fascicularis</i>) Induced by Mild Testicular Hyperthermia and Exogenous Testosterone Treatment	83
<i>Yue Jia, Amiya P. Sinha Hikim, Yan-He Lue, Ronald S. Swerdloff, Yanira Vera, Xue-Shen Zhang, Zhao-Yuan Hu, Yin-Chuan Li, Yi-Xun Liu, and Christina Wang</i> BCL2 phosphorylation and activation of the MAPK14-mediated mitochondria-dependent pathway are critical for male germ cell death in monkey.	

Regulation of Gene Expression in Bovine Blastocysts in Response to Oxygen and the Iron Chelator Desferrioxamine	93
<i>A.J. Harvey, K.L. Kind, and J.G. Thompson</i>	
Gene expression is regulated by hypoxia-inducible factors in bovine blastocysts, in response to reduced oxygen, or in the presence or absence of the iron chelator desferrioxamine.	
Gene Expression Profiles Linked to the Hormonal Induction of Male-Effect Pheromone Synthesis in Goats (<i>Capra hircus</i>)	102
<i>Yukihide Momozawa, Yukari Takeuchi, Masaru Kitago, Koji Masuda, Yoshie Kakuma, Chie Hashizume, Toru Ichimaru, Kazutaka Mogi, Hiroaki Okamura, Tomohiro Yonezawa, Takefumi Kikusui, and Yuji Mori</i>	
In parallel with male-effect pheromone biosynthesis, the expression of nine genes were found to increase; <i>SCD1</i> and <i>ELOVL5</i> genes increased 100-fold compared with pheromone-negative controls.	
Cysteine-Rich Secretory Protein 2 Binds to Mitogen-Activated Protein Kinase Kinase Kinase 11 in Mouse Sperm	108
<i>Gerard M. Gibbs, Deborah M. Bianco, Duangporn Jamsai, Amy Herlihy, Sika Ristevski, R. John Aitken, David M. de Kretser, and Moira K. O'Bryan</i>	
The C-terminal cysteine-rich domain of cysteine-rich secretory protein 2 (CRISP2) binds to the C-terminus of mitogen-activated protein kinase kinase kinase 11 (MAP3K11) present in the testis and epididymal sperm.	
A Novel Method to Isolate Primordial Germ Cells and Its Use for the Generation of Germline Chimeras in Chicken	115
<i>Yasuhiro Yamamoto, Fumitake Usui, Yoshiaki Nakamura, Yohei Ito, Takahiro Tagami, Keijiro Nirasawa, Yuko Matsubara, Tamao Ono, and Hiroshi Kagami</i>	
Functional chicken PGCs isolated by a novel method generated germline chimeras.	
Milk Caseins Decrease the Binding of the Major Bovine Seminal Plasma Proteins to Sperm and Prevent Lipid Loss from the Sperm Membrane During Sperm Storage	120
<i>Annick Bergeron, Yves Brindle, Patrick Blondin, and Puttaswamy Manjunath</i>	
Milk caseins prevent the detrimental effect of BSP proteins on the sperm membrane during sperm preservation.	
Identification, Isolation, and In Vitro Culture of Porcine Gonocytes	127
<i>Sandeep Goel, Miki Sugimoto, Naojiro Minami, Masayasu Yamada, Shinichi Kume, and Hiroshi Imai</i>	
The lectin <i>Dolichos biflorus</i> agglutinin (DBA) is a specific marker for porcine gonocytes and primitive spermatogonia, and isolated gonocytes can be cultured for 7 days without addition of any specific growth factors to the medium.	
Vitellogenin and Its Messenger RNA During Ovarian Development in the Female Blue Crab, <i>Callinectes sapidus</i>: Gene Expression, Synthesis, Transport, and Cleavage	138
<i>Nili Zmora, John Trant, Siu-Ming Chan, and J. Sook Chung</i>	
The temporal and tissue-specific expression, secretion, and uptake of vitellogenin along with the pathway for conversion to vitellin display some differences as compared to other crustaceans and may be related to the reproductive physiology of the female blue crab.	
Protein Kinase A-Independent cAMP Stimulation of Progesterone in a Luteal Cell Model Is Tyrosine Kinase Dependent but Phosphatidylinositol-3-Kinase and Mitogen-Activated Protein Kinase Independent	147
<i>Elie Needle, Kathy Piparo, Donna Cole, Carolyn Worrall, Ian Whitehead, Gwen Mahon, and Laura T. Goldsmith</i>	
cAMP-stimulated progesterone expression is PKA independent but tyrosine dependent.	
Vasoactive Peptides in the Luteolytic Process Activated by PGF2alpha in Pseudopregnant Rabbits at Different Luteal Stages	156
<i>Cristiano Boiti, Margherita Maranesi, Cecilia Dall'Aglio, Luisa Pascucci, Gabriele Breccia, Anna Gobbetti, and Massimo Zerani</i>	
PGF2alpha up-regulates the expression for the RAS/EDN1 system in CL of pseudopregnant rabbits, but only in midluteal phase does a reciprocal feedback mechanism with EDN1, mediated by ACE, induce luteolysis.	
Differential Gene Expression among the Proximal Segments of the Rat Epididymis Is Lost after Efferent Duct Ligation	165
<i>Terry T. Turner, Daniel S. Johnston, Joshua N. Finger, and Scott A. Jelinsky</i>	
Efferent duct ligation causes loss of differential gene expression by the proximal segments of the rat epididymis.	
Regulation of Expression of Fibroblast Growth Factor 7 in the Pig Uterus by Progesterone and Estradiol	172
<i>Hakhyun Ka, Saeed Al-Ramadan, David W. Erikson, Greg A. Johnson, Robert C. Burghardt, Thomas E. Spencer, Laurie A. Jaeger, and Fuller W. Bazer</i>	
Maximal expression of FGF7 in uterine LE requires progesterone-mediated down-regulation of progesterone receptor in LE, induction of prostaglandin(s) in uterine stroma, and direct effects of conceptus estrogens on LE.	
Changes in Lipids Containing Long- and Very Long-Chain Polyunsaturated Fatty Acids in Cryptorchid Rat Testes	181
<i>Natalia E. Furland, Eduardo N. Maldonado, Pablo Ayuza Aresti, and Marta I. Aveldaño</i>	
The rapid depletion of germ cells occurring in cryptorchidism was accompanied by a massive loss of glycerophospholipids and sphingomyelin and a substantial accumulation of triglycerides and cholesterol esters in the cells remaining in the testes.	

Contents by Category

Embryo

- 9 Impact of Dietary Fatty Acids on Oocyte Quality and Development in Lactating Dairy Cows. *Ali A. Fouladi-Nashta, Carlos G. Gutierrez, Jin G. Gong, Philip C. Garnsworthy, and Robert Webb*
- 93 Regulation of Gene Expression in Bovine Blastocysts in Response to Oxygen and the Iron Chelator Desferrioxamine. *A.J. Harvey, K.L. Kind, and J.G. Thompson*

Female Reproductive Tract

- 172 Regulation of Expression of Fibroblast Growth Factor 7 in the Pig Uterus by Progesterone and Estradiol. *Hakhyun Ka, Saeed Al-Ramadan, David W. Erikson, Greg A. Johnson, Robert C. Burghardt, Thomas E. Spencer, Laurie A. Jaeger, and Fuller W. Bazer*

Gamete Biology

- 2 Oxidative Metabolism of Pyruvate Is Required for Meiotic Maturation of Murine Oocytes In Vivo. *Mark T. Johnson, Edward A. Freeman, David K. Gardner, and Patricia A. Hunt*
- 37 *Catsper3* and *Catsper4* Are Essential for Sperm Hyperactivated Motility and Male Fertility in the Mouse. *Jingling Jin, Nange Jin, Huii Zheng, Seungil Ro, Dora Tafolla, Kenton M. Sanders, and Wei Yan*
- 71 Factors Maintaining Normal Sperm Tail Structure During Epididymal Maturation Studied in *Gopc^{-/-}* Mice. *Fumie Suzuki-Toyota, Chizuru Ito, Yoshiro Toyama, Mamiko Maekawa, Ryoji Yao, Tetsuo Noda, Hiroshi Iida, and Kiyotaka Toshimori*
- 108 Cysteine-Rich Secretory Protein 2 Binds to Mitogen-Activated Protein Kinase Kinase 11 in Mouse Sperm. *Gerard M. Gibbs, Deborah M. Bianco, Duangporn Jamsai, Amy Herlihy, Sika Ristevski, R. John Aitken, David M. de Kretser, and Moira K. O'Bryan*
- 115 A Novel Method to Isolate Primordial Germ Cells and Its Use for the Generation of Germline Chimeras in Chicken. *Yasuhiro Yamamoto, Fumitake Usui, Yoshiaki Nakamura, Yohei Ito, Takahiro Tagami, Keiji Nirasawa, Yuko Matsubara, Tamao Ono, and Hiroshi Kagami*

Male Reproductive Tract

- 165 Differential Gene Expression among the Proximal Segments of the Rat Epididymis Is Lost after Efferent Duct Ligation. *Terry T. Turner, Daniel S. Johnston, Joshua N. Finger, and Scott A. Jelinsky*

Mechanisms of Hormone Action

- 102 Gene Expression Profiles Linked to the Hormonal Induction of Male-Effect Pheromone Synthesis in Goats (*Capra hircus*). *Yukihide Momozawa, Yukari Takeuchi, Masaru Kitago, Koji Masuda, Yoshie Kakuma, Chie Hashizume, Toru Ichimaru, Kazutaka Mogi, Hiroaki Okamura, Tomohiro Yonezawa, Takefumi Kikusui, and Yuji Mori*

Ovary

- 18 Insulin-Like Growth Factor (IGF) 2 Stimulates Steroidogenesis and Mitosis of Bovine Granulosa Cells Through the IGF1 Receptor: Role of Follicle-Stimulating Hormone and IGF2 Receptor. *L.J. Spicer and P.Y. Aad*
- 28 Fibroblast Growth Factor 2 Is More Dynamic than Vascular Endothelial Growth Factor A During the Follicle-Luteal Transition in

the Cow. *R.S. Robinson, L.T. Nicklin, A.J. Hammond, D. Schams, M.G. Hunter, and G.E. Mann*

- 61 Actions of Epidermal Growth Factor Receptor/Mitogen-Activated Protein Kinase and Protein Kinase C Signaling in Granulosa Cells from *Gallus gallus* Are Dependent upon Stage of Differentiation. *Dori C. Woods, Morgan J. Haugen, and A.L. Johnson*
- 138 Vitellogenin and Its Messenger RNA During Ovarian Development in the Female Blue Crab, *Callinectes sapidus*: Gene Expression, Synthesis, Transport, and Cleavage. *Nili Zmora, John Trant, Siu-Ming Chan, and J. Sook Chung*
- 147 Protein Kinase A-Independent cAMP Stimulation of Progesterone in a Luteal Cell Model Is Tyrosine Kinase Dependent but Phosphatidylinositol-3-Kinase and Mitogen-Activated Protein Kinase Independent. *Elie Needle, Kathy Piparo, Donna Cole, Carolyn Worrall, Ian Whitehead, Gwen Mahon, and Laura T. Goldsmith*
- 156 Vasoactive Peptides in the Luteolytic Process Activated by PGF₂α in Pseudopregnant Rabbits at Different Luteal Stages. *Cristiano Boiti, Margherita Maranesi, Cecilia Dall'Aglio, Luisa Pascucci, Gabriele Brecchia, Anna Gobbetti, and Massimo Zerani*

Pregnancy

- 45 Modulation of Human Arterial Tone During Pregnancy: The Effect of the Bioactive Metabolite Sphingosine-1-Phosphate. *Nicola K. Hudson, Maureen O'Hara, Helen A. Lacey, Jemma Corcoran, Denise G. Hemmings, Mark Wareing, Philip Baker, and Michael J. Taggart*

Reproductive Technology

- 120 Milk Caseins Decrease the Binding of the Major Bovine Seminal Plasma Proteins to Sperm and Prevent Lipid Loss from the Sperm Membrane During Sperm Storage. *Annick Bergeron, Yves Brindle, Patrick Blondin, and Puttaswamy Manjunath*

Testis

- 83 Signaling Pathways for Germ Cell Death in Adult Cynomolgus Monkeys (*Macaca fascicularis*) Induced by Mild Testicular Hyperthermia and Exogenous Testosterone Treatment. *Yue Jia, Amiya P. Sinha Hikim, Yan-He Lue, Ronald S. Swerdloff, Yanira Vera, Xue-Shen Zhang, Zhao-Yuan Hu, Yin-Chuan Li, Yi-Xun Liu, and Christina Wang*
- 127 Identification, Isolation, and In Vitro Culture of Porcine Gonocytes. *Sandeep Goel, Miki Sugimoto, Naojiro Minami, Masayasu Yamada, Shinichi Kume, and Hiroshi Imai*
- 181 Changes in Lipids Containing Long- and Very Long-Chain Polyunsaturated Fatty Acids in Cryptorchid Rat Testes. *Natalia E. Furland, Eduardo N. Maldonado, Pablo Ayuza Aresti, and Marta I. Aveldaño*

Toxicology

- 53 Epidermal Growth Factor-Like Growth Factors Prevent Apoptosis of Alcohol-Exposed Human Placental Cytotrophoblast Cells. *Garen S. Wolff, Po Jen Chiang, Susan M. Smith, Roberto Romero, and D. Randall Armant*