

BIOLOGY *of* REPRODUCTION

Official Journal of the Society for
the Study of Reproduction

AUGUST 2007

VOLUME 77 NUMBER 2

Biology of Reproduction Highlights	189
Minireview. Polyunsaturated Fatty Acids in Male and Female Reproduction.	190
<i>D. Claire Wathes, D. Robert E. Abayasekara, and R. John Aitken</i>	
Polyunsaturated fatty acids (PUFAs) can affect reproduction through changes to prostaglandin and steroid synthesis and membrane properties of the gametes, but conclusive evidence that dietary consumption of particular PUFAs improves aspects of fertility is lacking.	
Relationships of Follicle Versus Oocyte Maturity to Ultrasound Morphology, Blood Flow, and Hormone Concentrations of the Preovulatory Follicle in Mares	202
<i>O.J. Ginther, E.L. Gastal, M.O. Gastal, M.A.R. Siddiqui, and M.A. Beg</i>	
Follicular-fluid estradiol, progesterone, and the IGF system are associated with final oocyte maturation but not with follicular maturation.	
Constitutive Steroidogenesis in Ovine Large Luteal Cells May Be Mediated by Tonically Active Protein Kinase A	209
<i>Randy L. Bogan and Gordon D. Niswender</i>	
A novel assay that quantifies concentrations and relative phosphorylation states of steroidogenic acute regulatory protein provided evidence that constitutive steroidogenesis in ovine large luteal cells is mediated by tonically active PKA.	
Impaired Progesterone Production in <i>Nr5a2</i>^{+/-} Mice Leads to a Reduction in Female Reproductive Function	217
<i>Cassandre Labelle-Dumais, Jean-François Paré, Luc Bélanger, Riaz Farookhi, and Daniel Dufort</i>	
The nuclear receptor NR5A2 is involved in progesterone production and female reproductive function in vivo in the mouse.	
The Hedgehog Signaling Pathway in the Mouse Ovary	226
<i>Matthew C. Russell, Robert G. Cowan, Rebecca M. Harman, Ashleigh L. Walker, and Susan M. Quirk</i>	
Components of the hedgehog cell signaling pathway are expressed in the mouse ovary and modulate granulosa cell proliferation and steroidogenesis.	
Genetic Factors Contributing to Defective Spermatogonial Differentiation in Juvenile Spermatogonial Depletion (<i>Utp14b</i>^{jsd}) Mice	237
<i>Olga U. Bolden-Tiller, Helio Chiarini-Garcia, Christophe Poirier, Daniel Alves-Freitas, Connie C. Weng, Gunapala Shetty, and Marvin L. Meistrich</i>	
The genetic background, particularly the Y chromosome, affects the rate and severity of spermatogonial depletion and the restoration of spermatogenesis following the suppression of testosterone or cryptorchidization in <i>Utp14b</i> ^{jsd} mice.	
Depletion of Ovarian Follicles with Age in Chimpanzees: Similarities to Humans.	247
<i>K.P. Jones, L.C. Walker, D. Anderson, A. Lacreuse, S.L. Robson, and K. Hawkes</i>	
The rate of decline in primordial follicle numbers in chimpanzee ovaries is very similar to the previously reported rate of decline in human ovaries over the same range of ages.	
Differential Effects of Various Estradiol-17beta Treatments on Follicle-Stimulating Hormone Peaks, Luteinizing Hormone Pulses, Basal Gonadotropin Concentrations, and Antral Follicle and Luteal Development in Cyclic Ewes	252
<i>David M.W. Barrett, Rajesha Duggavathi, Kate L. Davies, Pawel M. Bartlewski, Edward T. Bagu, and Norman C. Rawlings</i>	
Various estradiol-17beta concentrations and exposure times altered antral follicle development and differentially effected FSH peaks, LH pulses, basal gonadotropin concentrations, and CL function.	
Role of PI3-Kinase and PI4-Kinase in Actin Polymerization During Bovine Sperm Capacitation	263
<i>Nir Etkovitz, Sara Rubinstein, Limor Daniel, and Haim Breitbart</i>	
Actin polymerization during sperm capacitation depends on PI4-kinase activity and is independent of PI3-kinase activity.	
An Impaired Breeding Phenotype in Mice with a Genetic Deletion of Beta-2 Microglobulin and Diminished MHC Class I Expression: Role in Reproductive Fitness	274
<i>Joanne C. Cooper, Gillian B. Dealtry, Mohamed Abdelrahman Ahmed, Petra C. Arck, Burghard F. Klapp, Sandra M. Blois, and Nelson Fernández</i>	
A selective deletion of the gene encoding B2M, a protein that associates with the heavy chain of the MHC Class I products, decreases reproductive fitness of C57BL6 mice by affecting breeding performance, litter size, and weaning success.	

Reproductive-Tactic-Specific Variation in Sperm Swimming Speeds in a Shell-Brooding Cichlid	280
<i>J.L. Fitzpatrick, J.K. Desjardins, N. Milligan, R. Montgomerie, and S. Balshine</i>	
In a fish species characterized by four alternative male reproductive tactics, sperm from males performing sneak fertilizations swim faster than sperm from males adopting other reproductive tactics.	
Buffalos (<i>Bubalus bubalis</i>) Cloned by Nuclear Transfer of Somatic Cells	285
<i>Deshun Shi, Fenghua Lu, Yingming Wei, Kuiqing Cui, Sufang Yang, Jingwei Wei, and Qingyou Liu</i>	
Three cloned buffalos from fetal fibroblasts and adult ovarian granulosa cells were produced.	
Pig Conceptuses Increase Uterine Interferon-Regulatory Factor 1 (<i>IRF1</i>), but Restrict Expression to Stroma Through Estrogen-Induced <i>IRF2</i> in Luminal Epithelium	292
<i>Margaret M. Joyce, James R. Burghardt, Robert C. Burghardt, R. Neil Hooper, Laurie A. Jaeger, Thomas E. Spencer, Fuller W. Bazer, and Greg A. Johnson</i>	
Pig conceptuses secrete estrogens and interferons to differentially regulate <i>IRF1</i> and <i>IRF2</i> in endometrial stroma and luminal epithelium.	
Chronic Cyclophosphamide Exposure Alters the Profile of Rat Sperm Nuclear Matrix Proteins	303
<i>Alexis M. Codrington, Barbara F. Hales, and Bernard Robaire</i>	
The expression of several spermatozoal nuclear matrix protein components, a number of which have been identified for the first time, is altered following exposure to the alkylating agent and male-mediated developmental toxicant, cyclophosphamide.	
Microarray Analyses of Newborn Mouse Ovaries Lacking <i>Nobox</i>	312
<i>Youngsok Choi, Yingying Qin, Michael F. Berger, Daniel J. Ballow, Martha L. Bulyk, and Aleksandar Rajkovic</i>	
<i>Nobox</i> activates expression of oocyte-specific genes and represses transcription of genes preferentially expressed in testes.	
Differential Modulation of Gonadotropin Secretion by Selective Estrogen Receptor 1 and Estrogen Receptor 2 Agonists in Ovariectomized Ewes	320
<i>J. Alejandro Arreguin-Arevalo, Tracy L. Davis, and Terry M. Nett</i>	
The positive and negative feedback of estradiol on LH secretion is mediated by <i>ESR1</i> and <i>ESR2</i> in OVX ewes.	
Expression Analysis of the Mouse Multi-Copy X-Linked Gene <i>Xlr-Related, Meiosis-Regulated (Xmr)</i>, Reveals That <i>Xmr</i> Encodes a Spermatid-Expressed Cytoplasmic Protein, <i>SLX/XMR</i>	329
<i>Louise N. Reynard, James M.A. Turner, Julie Cocquet, Shantha K. Mahadevaiah, Aminata Touré, Christer Höög, and Paul S. Burgoyne</i>	
<i>XMR/SLX</i> protein is a spermatid-specific cytoplasmic protein.	
Evaluation of Chromosomal Risk Following Intracytoplasmic Sperm Injection in the Mouse	336
<i>Hiroyuki Tateno and Yujiroh Kamiguchi</i>	
The incidence of ICSI embryos with structural chromosome aberrations is dependent on the kind of medium used for sperm incubation and the time for sperm incubation.	
Maternal and Fetal Growth, Body Composition, Endocrinology, and Metabolic Status in Undernourished Adolescent Sheep	343
<i>Justin Luther, Raymond Aitken, John Milne, Masatoshi Matsuzaki, Lawrence Reynolds, Dale Redmer, and Jacqueline Wallace</i>	
Limiting maternal intake during adolescent pregnancy gradually depleted maternal body reserves, impaired fetal nutrient supply, and slowed fetal soft tissue growth.	
Placental Growth, Angiogenic Gene Expression, and Vascular Development in Undernourished Adolescent Sheep	351
<i>Justin Luther, John Milne, Raymond Aitken, Masatoshi Matsuzaki, Lawrence Reynolds, Dale Redmer, and Jacqueline Wallace</i>	
Limiting maternal intake during adolescent pregnancy does not influence placental mass per se but impairs vascular development in the maternal caruncle, which may in turn impact fetal nutrient supply.	
Wild Fulvous Fruit Bats (<i>Rousettus leschenaulti</i>) Exhibit Human-Like Menstrual Cycle	358
<i>Xiaoping Zhang, Cheng Zhu, Haiyan Lin, Qing Yang, Qizhi Ou, Yuchun Li, Zhong Chen, Paul Racey, Shuyi Zhang, and Hongmei Wang</i>	
Wild fulvous fruit bats menstruate in a similar way to humans and may be useful non-primate animal models to study menstruation and menstrual dysfunction.	
The Role of Aquaporin 3 in the Movement of Water and Cryoprotectants in Mouse Morulae	365
<i>Keisuke Edashige, Satoshi Ohta, Mitsunobu Tanaka, Tatsunaga Kuwano, Delgado M. Valdez, Jr., Takao Hara, Bo Jin, Sei-ichi Takahashi, Shinsuke Seki, Chihiro Koshimoto, and Magosaburo Kasai</i>	
Water and cryoprotectants move through aquaporin 3 in mouse morulae.	
Sperm Motility Defects and Infertility in Male Mice with a Mutation in <i>Nsun7</i>, a Member of the Sun Domain-Containing Family of Putative RNA Methyltransferases	376
<i>Tanya Harris, Becky Marquez, Susan Suarez, and John Schimenti</i>	
An ENU-induced male infertility mutation in mice was traced to <i>Nsun7</i> , which encodes a putative RNA methyltransferase, resulting in decreased progressive motility and a midpiece mitochondrial defect.	
Position Announcement and Reviews Editor Announcement	
SSR 2008 and the World Congress on Reproductive Biology	