

LEPTIN ENHANCES BOVINE ADENOHYPOPHYSEAL LH RELEASE

Leptin Acts at the Bovine Adenohypophysis to Enhance Basal and GnRH-Mediated Release of LH: Differential Effects Are Dependent Upon Nutritional History¹

5

M. Amstalden,^{3,4,5} D. A. Zieba,^{3,4,5} J. F. Edwards,⁶ P.G. Harms,⁴ T. H. Welsh, Jr.,⁴ R. L. Stanko,^{3,7} G. L. Williams^{2,3,4,5}

Animal Reproduction Laboratory,³ Texas A&M University Agricultural Research Station,

Beeville, Texas 78102; Department of Animal Science⁴ and Center for Animal Biotechnology

10 *and Genomics⁵, Texas A&M University, College Station, Texas 77843; Department of*

Pathobiology⁶, College of Veterinary Medicine, Texas A&M University, College Station, Texas

77843; Department of Animal and Wildlife Sciences⁷, Texas A&M University-Kingsville,

Kingsville, Texas 78363

Received:

15 First decision:

Accepted:

¹Supported by USDA-NRI 00-35203-9132. A preliminary report of this work was presented at the 35th Annual Meeting to the Society for the Study of Reproduction.

²Correspondence: G.L. Williams, Animal Reproduction Laboratory, Texas A&M University

20 Agricultural Research Station, 3507 Hwy 59 E, Beeville, TX 78102-9410; FAX 361 358 4930;

e-mail: glw@fnbnet.net

Key words: hypothalamus, anterior pituitary, leptin, luteinizing hormone, gonadotropin-releasing hormone

ABSTRACT

25 Recombinant ovine leptin (oleptin) stimulates an acute increase in the secretion of LH in
fasted, but not in normal-fed, cows through an augmentation of the magnitude of individual
pulses of LH. Herein, we tested the hypothesis that this effect could be accounted for by
functional changes at the adenohipophyseal level. Eleven ovariectomized, estradiol-implanted
cows were assigned to one of two dietary groups: 1) Normal-Fed (n=6) and 2) Fasted (fasted for
30 72 h; n=5). Adenohipophyseal (AP) explants collected following euthanasia were perfused with
Krebs-Ringer bicarbonate buffer (KRB) for a total of 6.5 h, including a 2-h treatment at 2.5 h
with KRB or increasing doses of oleptin, and a challenge with 50 ng of GnRH at 4.5 h. To test
for effects of leptin at the hypothalamic level, explants encompassing the medial basal
hypothalamus-infundibular complex (HYP) were incubated in KRB alone (control) or KRB
35 containing 1000 ng of oleptin. Basal release of LH from AP explants treated with leptin was
greater ($P < 0.02$) than control-treated explants in fasted but not in normal-fed cows. To the
contrary, leptin-treated explants from normal-fed, but not fasted, cows released more ($P < 0.001$)
LH in response to GnRH than control-treated tissues. Neither fasting nor leptin affected ($P >$
0.1) the secretion of GnRH from HYP explants. These observations support the hypothesis that
40 leptin modulates the secretion of LH in mature cows to a large extent by its direct actions at the
adenohipophysis. Differential manifestations of these effects are dependent upon nutritional
history.

INTRODUCTION

45 Leptin, a 16 kDa product of the *ob* gene, plays a major role in communicating nutritional
status to the central nervous system, including centers that control reproduction [1-4]. The leptin
receptor (LR) is present at both hypothalamic and adenohipophyseal loci [5-8], and in vitro

studies using explants collected from normal-fed rodents indicate that leptin can act directly at both sites [9, 10] to stimulate the release of GnRH and LH, respectively. However, in domestic ruminants, leptin has been shown repeatedly to stimulate LH secretion only during periods of negative energy balance. **In addition, leptin receptor mRNA increases in feed-restricted ewes [7].** In this context, restriction of nutrient intake **for 2-3 d** reduces concentrations of leptin concurrent with reductions in the frequency of LH pulses in castrated male sheep [11], and peripubertal heifers [12]. Leptin treatment prevents these reductions, suggesting an effect at hypothalamic centers that regulate pulsatile GnRH secretion [11, 13].

Importantly, leptin also stimulates an increase in LH secretion in animals in which short-term restrictions in nutrient intake does not measurably alter the pattern of LH release. For example, **2-3 d of** fasting reduces expression of the leptin gene and markedly diminishes circulating concentrations of leptin, insulin, and IGF-1 in mature ruminants, without reducing mean concentrations of LH or the frequency of LH pulses [11, 14-16]. Nonetheless, 2-3 d of total feed restriction clearly hypersensitizes the mature cow to leptin, resulting in acute increases in the mean baseline and the magnitude of individual pulses of LH [16, 17]. This would suggest that the primary effects of leptin in this animal model are at the adenohipophyseal level. To test this hypothesis further, we examined in vitro the stimulatory effects of leptin on basal and GnRH-mediated secretion of LH from adenohipophyseal explants obtained from fasted and non-fasted cows. To assess coincident hypothalamic responses, the effects of leptin on GnRH secretion from hypothalamic-infundibular explants were also examined.

MATERIALS AND METHODS

All animal-related procedures used in these studies were approved by the Institutional Agricultural Animal Care and Use Committee of The Texas A&M University System.

Animal Model and Procedures

Eleven mature, ovariectomized cows, each bearing an estradiol implant to maintain circulating estradiol at 2-5 pg/ml, were used. This animal model has been used extensively by our laboratory and others [16-19] to study the effects of nutrition on neuroendocrine control of gonadotropin secretion in cattle. With this approach, implants provide a constant level of steroid (estradiol) negative feedback without the complications associated with ovarian cyclicity. Cows used in the current experiments had mean (\pm SEM) concentrations of estradiol of 4.35 ± 0.45 pg/ml.

Cows, in moderately thin body condition (BCS=4; 1-9 scale), were fed once daily at 0700 h a diet formulated to provide 100% of the National Research Council (NRC) [20] requirements for maintenance for at least 2 weeks before the beginning of the experiment. Each cow was assigned randomly to one of two dietary groups: 1) Normal Fed; cows were fed 100 % of the NRC [20] requirements to maintain body weight (n=6), and 2) Fasted; cows were fasted for 72 h with free access to water (n=5). On the day before the start of dietary treatments (Day – 1), cows were fitted with jugular catheters (polyethylene tubing, 1.4 mm inside diameter, 1.9 mm outside diameter; Becton Dickinson, Parsippany, NJ) for intensive blood sampling. Cows were placed in stanchions and blood was collected at 10-min intervals for 6 h on Days 0 and 3 of the experiment via an extension connected to the jugular catheter. Blood samples were dispensed into tubes containing 150 μ l of a solution containing heparin (1000 IU/ml) and 5% EDTA and placed immediately on ice. Plasma was harvested by centrifugation and stored at -20°C until hormone analysis. At the end of the intensive blood sampling on Day 3, cows were humanely euthanized by exsanguination following captive bolt stunning. Diencephalons were removed

after disconnection of infundibuli from adenohypophyses. Adenohypophyses were removed from the sella turcica and kept on ice until tissue processing.

95 ***Effects of Leptin on Basal and GnRH-Mediated Release of LH from AP Explants***

Adenohypophyses collected from each cow were dissected and sliced sagittally into approximately 0.5 x 2 mm strips. Four adenohypophyseal (AP) strips **from each adenohypophysis** were selected randomly, placed into each of 5 perfusion chambers, and perfused with Krebs-Ringer bicarbonate buffer (KRB) for 6.5 h using a multiple micro chamber
100 perfusion system (Endotronics Inc., Coon Rapids, MN). **Thus, adenohypophyseal explants from each cow were perfused with each dose of leptin tested or control media.** In preliminary studies, it was determined that basal release of LH by AP explants was attained within the first 2.5 h of perfusion. At 2.5 h, AP explants were perfused for 2 h with KRB containing 0 (control), 5, 10, 50, or 100 ng/ml of oleptin. After 4.5 h of perfusion, AP explants
105 were challenged with 50 ng of GnRH (Sigma, St. Louis, MO) via an injection port in the chamber (Fig. 1A). **In addition to investigating the effects of leptin on GnRH-mediated release of LH, the stimulation of AP explants by GnRH also made it possible to verify the viability of tissues based on specific stimulation of gonadotropin release.**

Approximately 1-ml fractions of perfused media were collected at 10-min intervals and
110 stored at -20° C until analysis for LH. In a separate study, we have reported simultaneous effects of leptin on basal and growth hormone (GH)-releasing hormone-mediated GH release from these same explants [21].

Effects of Leptin on GnRH Secretion from the Isolated Medial Basal Hypothalamus-Infundibular Complex

115 To isolate the medial basal hypothalamus (MBH), the entire hypothalamic-infundibular
complex was transected sagittally into equal halves. A cut was made 2 mm lateral to the median
sagittal cut for each half, followed by an anterior coronal cut extending from the optic chiasm to
the anterior commissure. A posterior coronal cut was made resulting in a section containing
approximately half of the mamillary body. A final transverse cut containing two-thirds of the
120 mamillary body was performed. Each half of the medial basal hypothalamus connected to its
infundibular half was incubated in 0.7 ml KRB for 5 h. Medium was collected and replaced
every 30 min. After a 2.5 h period of equilibration, hypothalamic-infundibular (HYP) explants
from each cow were incubated with KRB containing either 0 or 1000 ng of oleptin for 1 h, so
that one-half served as a control for the corresponding half (Fig. 1B). At 4 h of incubation, all
125 explants were incubated with KRB containing 60 mM K⁺ to test tissue viability, based on
nonspecific stimulation of GnRH release [22].

Doses of recombinant oleptin used to treat HYP explants were chosen empirically based
on 1) quantities shown to have positive effects in comparable tissue collected from rodents [9]
and 2) preliminary experiments using tissues collected from bulls and steers at slaughter. In
130 these preliminary experiments, GnRH released from HYP explants collected from full-fed cattle
was not affected by leptin (1 to 1000 ng of recombinant oleptin for 30 to 60 min). This
observation was not totally surprising, since the stimulatory effects of leptin on LH release
observed in vivo [11] were limited to fasted cows. Thus, we chose to use 1000 ng of oleptin in
the current experiment.

135 *RIAs*

Concentrations of GnRH in HYP culture media were determined by a validated assay as
described previously [22]. Plasma concentrations of LH and concentrations of LH in AP

perifused media were determined as reported previously [23]. Intra- and inter-assay coefficients of variation for these assays averaged 6.2 and 10.7 %, respectively. Serum estradiol was assayed
140 in extracted samples in a single assay as reported previously [24].

Statistical Analysis

Circulating concentrations of LH were analyzed by general linear mixed models for repeated measures using the Mixed procedure (PROC MIXED) of the Statistical Analysis System (SAS 8.1; SAS Institute Inc., Cary NC). The frequency and amplitude of LH pulses
145 were determined using both visual inspection and a pulse-detection algorithm, Pulsefit 1.2 [25]. Sources of variation were diet, day, and diet x day. Day was used as the repeated variable and cow(diet) was used as the subject.

To test the effects of diet, leptin, and GnRH treatments on the release of LH from AP explants in vitro, the 6.5-h period of perfusion was subdivided in three periods (I-III),
150 corresponding to Equilibration (0 to 2.5 h), Basal Release (2.5 to 4.5 h), and GnRH Stimulation (4.5 to 6.5 h), respectively. Hormone data obtained from perifused media were analyzed by the general linear mixed model (PROC MIXED) procedure of SAS. Sources of variation were diet, leptin treatment, period, and all interactions. Period was used as the repeated variable and cow(diet) was used as subject. The Least Squares means procedure was used to compare means
155 when a significant F-value was obtained. Because of differences in the release of LH by AP explants among perfusion chambers at the end of the Equilibration period, covariate analyses were performed to test main effects during Period II (Basal Release). Mean concentrations of LH in the last 3 samples collected during equilibration were used as the covariate (Covariate 1). To test main effects during Period III (GnRH Stimulation), another covariate analysis was
160 performed, using the mean of the last 3 samples of Period II (Basal) as covariate (Covariate 2).

To test cumulative (Basal + GnRH) effects of leptin and GnRH-stimulation on the amount of LH released by AP explants, an analysis was performed using Covariate 1 to test main effects during Period III.

To test the effects of leptin on GnRH release from HYP explants, data were analyzed by the general linear mixed model procedure of SAS (PROC MIXED). Sources of variation were diet, leptin treatment, period, and all possible interactions. Period was used as the repeated variable and cow(diet) was used as subject.

RESULTS

Does Leptin Modulate Basal or GnRH-Mediated LH release from the Adenohypophysis?

As expected, mean plasma concentrations and mean frequency and amplitude of LH pulses did not differ ($P > 0.1$) between Fasted and Normal-Fed groups before euthanasia (Table 1). Using covariate analysis to adjust for differences in basal release of LH among perfusion chambers, it was observed that dietary treatment interacted with in vitro leptin treatments to influence the release of LH from AP explants. Leptin-treated AP explants from fasted cows had a higher ($P < 0.02$) basal release of LH than control-treated explants. Because effects of individual doses of leptin on the basal release of LH from AP explants of fasted cows did not differ, all doses of leptin were combined for presentation (Fig. 2). In explants of normal-fed cows, leptin did not affect ($P > 0.1$) basal release of LH (Fig. 2).

In contrast to observations made for basal release, all doses of leptin increased ($P < 0.001$) GnRH-mediated release of LH from AP explants of normal-fed cows, but not fasted cows (Fig. 3). In fact, in the Fasted group, AP explants treated with 5 ng/ml of oleptin released less ($P < 0.001$; 1059 ± 55.4 ng/ml) LH in response to GnRH than control-treated (1133 ± 62.6 ng/ml) explants, and all other doses had no effect on this variable. Since leptin increased basal release

of LH from AP explants obtained from fasted cows before GnRH treatment, followed by non-
185 significant numerical increases after GnRH, we examined the combined effects of leptin on basal
and GnRH-stimulated release of LH. Leptin at doses of 50 (1416 ± 87.6 ng/ml) and 100 ng/ml
(1455 ± 89.1 ng/ml) increased ($P < 0.001$) the overall (basal plus GnRH-stimulated) release of
LH compared to control-treated explants.

Effects of Leptin on the Secretion of GnRH from the Medial Basal Hypothalamus-

190 *Infundibular Complex*

Data from one normal-fed cow were not included in the analysis due to a lack of response
to 60 mM K^+ , suggesting a loss of tissue viability. Neither fasting nor leptin affected ($P > 0.1$)
basal or potassium-stimulated release of GnRH from HYP explants (Fig. 4).

DISCUSSION

195 In the current study, we examined the hypothesis that the hypersecretion of LH induced
by leptin in vivo in fasted, mature cows [16] could be explained by effects at the
adenohypophyseal level. Significant effects of leptin on basal release of LH were observed in
AP explants from fasted cows, on AP responsiveness to GnRH in normal-fed cows, and on
overall release (basal plus GnRH-stimulated release of LH) in AP explants of fasted cows
200 perfused with the highest doses of leptin (50 and 100 ng/ml). In contrast, we were unable to
detect effects of leptin on the release of GnRH from isolated HYP explants obtained from either
normal-fed or fasted cows.

As noted previously, **2-3 d of** restriction of feed intake does not affect mean
concentrations of LH or its secretory patterns in mature ewes or cows [11, 14 – 16], although
205 signs of metabolic stress, including decreased circulating insulin, IGF-I, and leptin, are clearly
present. Similar metabolic constraints are observed in fasted heifers [12, 13] and estradiol-

implanted wethers [11], but in these ruminant animal models, short-term fasting is able to reduce the frequency of LH pulses similar to that observed for monogastric species [26 – 28].

Nonetheless, feed-restricted, mature cows are responsive to leptin treatment and mean concentrations, mean baseline, and the magnitude of pulses of LH are increased [16]. Questions then arise as to whether stimulation of LH release by leptin is accounted for by effects at hypothalamic and (or) adenohipophyseal levels. Most studies have proposed that leptin stimulates secretion of LH via interaction with neuronal systems and ultimately, stimulation of GnRH secretory neurons. It is likely that, in models in which the frequency of LH pulses is diminished by feed restriction [11, 13], leptin stimulates GnRH release to prevent the effects of fasting. Alternatively, diminished release of LH during fasting could be explained by changes in the amplitude of GnRH pulses and its ability to trigger LH release, as observed previously in this laboratory [18] and others [29] in other physiological contexts. However, under feed-restricted conditions, in which frequency of LH pulses is not affected, as observed in mature female ruminants [16], the stimulatory effects of leptin on LH secretion are more plausibly explained by a direct action at the adenohipophysis. In addition to the current results, evidence supporting a direct action of leptin at the level of the adenohipophysis includes the localization of leptin receptors on gonadotropes of ewes [8] and leptin stimulation of LH release from rat anterior pituitaries in vitro [9]. However, in contrast to the study performed using rat tissue [9], we observed no effect of leptin on basal release of LH from AP explants from normal-fed cows, suggesting important species differences in the acute interaction of nutritional status and leptin with gonadotropes in cattle. This view is supported further by our observation that LH release from primary cultures of bovine adenohipophyseal cells, maintained under optimal nutrient

conditions, was not affected by leptin at doses of 1 to 1000 ng/ml (Amstalden et al., unpublished
230 observations).

Increases in basal release of LH from AP explants from fasted cows treated with leptin
appear differently than what would be expected from a classical secretagogue. In fact, after the
period of equilibration (initial 2.5 h of perfusion) and before GnRH stimulation (last 2 h of
perfusion), basal release of LH tended to continue to decrease in all treatments, independent of
235 dietary group. **This steady decline in the release of LH during the Basal-Release period may
reflect the absence of stimulation by endogenous GnRH.** However, the decrease in LH
release by AP explants from fasted cows treated with leptin was significantly smaller than
control-treated explants. This effect was not observed in explants from normal-fed cows. It is
possible that leptin may have stimulated an increase in basal metabolism of gonadotropes from
240 fasted cows, resulting in an elevated release of LH compared to gonadotropes of control-treated
explants. In support of this concept, increased glucose uptake has been observed in muscle and
adipose tissue treated with leptin [30].

Mechanisms involved in the leptin-mediated increase in responsiveness to GnRH of AP
explants from normal-fed cows are unknown. However, there are several potential pathways
245 through which these effects could occur, including effects on Ca^{2+} ion channels, an increase in
the releasable pool of LH, and/or GnRH receptor desensitization. All of these are important for
cellular exocytosis of gonadotropins [31]. Leptin receptor signaling involves activation of Janus
kinase (JAK) and signal transducers and activators of transcription (STAT) pathways [32]. In
addition, mitogen-activated protein kinase [32] and phosphatidylinositol-3 kinase [33] have also
250 been involved in leptin receptor signaling. In porcine chromaffin cells, leptin caused a sustained
increase of intracellular Ca^{2+} and activated inositol 1,4,5-triphosphate production [34],

intracellular factors known to be associated with GnRH-receptor signaling and release of LH [35].

255 The lack of increased responsiveness to GnRH in AP explants from fasted cows treated with leptin is unclear. However, it is well established that GnRH stimulation of gonadotropes is necessary for synthesis of gonadotropins [36]. Thus, it is possible that the increased basal release of LH observed in leptin-treated explants of fasted cows and lack of any GnRH stimulation during this period caused a diminution of readily releasable pools of LH that were not replenished.

260 Experimental models describing the temporal secretion pattern of GnRH by the hypothalamus have been developed in several large species. In sheep, this has been accomplished through collection of hypophyseal-portal blood [37]. In the mature cow, the anatomy of the cranium impairs practical application of this technique. However, collection of cerebrospinal fluid from the third ventricle has been shown by our laboratory and others to be a
265 useful approach for determining patterns of GnRH secretion in cattle and sheep [18, 37], and studies are currently underway to investigate the effects of leptin using this approach. Incubation of HYP explants has also been used extensively to study the neuroendocrine regulation of GnRH release [38 - 40] in several species. In vitro incubations of isolated HYP tissue allows the investigation of factors acting directly at the hypothalamus. Hypothalamic explants used in the
270 current experiment included the MBH connected to the infundibulum. The arcuate nucleus, which rests within the MBH, contains leptin receptors [7] as well as neuronal cell bodies that express mediators of the effects of leptin within the central nervous system, such as neuropeptide Y, agouti-related peptide, and POMC (precursor of α -melanocyte-stimulating hormone) [1]. Thus, explants used in the current experiment contain at least part of the neuronal systems that

275 potentially mediate the effects of leptin on GnRH neurons. Using this approach, we were unable to detect effects of leptin on GnRH secretion from hypothalamus harvested from either normal-fed or fasted cows. In contrast, Yu et al. [9] have reported that leptin stimulates GnRH release from rat hypothalamic explants. It is not clear whether the discordant results are due to species differences or experimental methodology used to detect leptin effects.

280 Mechanisms that explain the inability of leptin to independently stimulate an increase in circulating concentrations of LH in cattle and sheep that are fed either maintenance [11, 41, 42] or growing diets [43] are not understood. However, the inability of leptin to stimulate directly basal secretion of LH in AP explants of normal-fed cows in the current study supports those findings. Syndromes involving the development of resistance to leptin have been observed in
285 obese humans [44, 45] and rodents [46, 47], and there is evidence that excessive activity of suppressor of cytokine signaling (SOCS)-3 activity is involved in leptin resistance [48, 49]. Inhibition of leptin signaling by SOCS-3 seems to involve suppression of JAK2 activation [48]. Whether a leptin-resistant condition occurs in ruminants that are under neutral or positive energy balance has not been determined; however, leptin is able to decrease feed intake in sheep [41,
290 42]. Moreover, the mass of ligand reaching target tissues must also be considered. We have reported recently that intravenously-injected recombinant oleptin causes an inverse, dose-related increase in basal plasma concentrations of LH in ovariectomized, estradiol-implanted cows fasted for 60 h [17]. A dose of 0.2 $\mu\text{g}/\text{kg}$ maximized the increase in LH, whereas doses of 2 and 20 $\mu\text{g}/\text{kg}$ caused lower and no responses, respectively. Reports in rodents [9] confirm that AP
295 explants treated with high concentrations of leptin (10^{-5} M) lose their ability to respond with an increase in LH release. Therefore, the duration and/or amount of exposure to the hormone likely determines the development of resistance.

In conclusion, results from the current experiment support the hypothesis that leptin affects the secretion of LH in the mature cow, in part, by its direct action at the adenohypophysis.

300 Leptin maintained higher basal AP release of LH in tissues obtained only from fasted cows, whereas increased responsiveness to GnRH was evident in AP explants obtained from normal-fed cows. This indicates that, while leptin acts on gonadotrope function under both normal and fasting conditions, the manifestations of these effects are expressed differentially and will require further study at the cellular level to characterize in detail.

305 **ACKNOWLEDGMENTS**

We acknowledge the National Pituitary Hormone Program for providing LH preparations, Dr. Jerry Reeves for the LH antisera, Dr. Terry Nett for the GnRH antisera, and Dr. Duane Keisler for recombinant leptin. We also acknowledge with gratitude the careful animal care and assistance of Melvin Davis and Randle Franke, and the technical assistance of Pat Chen,
310 Joanna Gallino, Marsha Green, and Stephanie Morton.

REFERENCES

1. Ahima RS, Saper CB, Flier JS, Elmquist JK. Leptin regulation of neuroendocrine systems. *Front Neuroendocrinol* 2000; 21:263-307.
2. Cunningham MJ, Clifton DK, Steiner RA. Leptin's action on the reproductive axis: perspectives and mechanisms. *Biol Reprod* 1999; 60:216-222.
3. Smith GD, Jackson LM, Foster DL. Leptin regulation of reproductive function and fertility. *Theriogenology* 2002; 57:73-86.
4. Williams GL, Amstalden M, Garcia MR, Stanko RL, Nizielski SE, Morrison CD, Keisler DH. Leptin and its role in the central regulation of reproduction in cattle. *Domest Anim Endo* 2002; 23:339-349.

5. Zamorano PL, Mahesh VB, De Sevilla LM, Chorich LP, Bhat GK, Brann DW. Expression and localization of the leptin receptor in endocrine and neuroendocrine tissues of the rat. *Neuroendocrinology* 1997; 65:223-228.
6. Lin J, Barb CR, Matteri RL, Kraeling RR, Chen X, Meinersmann RJ, Rampacek GB. Long form leptin receptor mRNA expression in the brain, pituitary, and other tissues in the pig. *Domest Anim Endocrinol* 2001; 19:53-61.
7. Dyer CJ, Simmons JM, Matteri RL, Keisler DH. Leptin receptor mRNA is expressed in ewe anterior pituitary and adipose tissues and is differentially expressed in hypothalamic regions of well-fed and feed-restricted ewes. *Domest Anim Endocrinol* 1997; 14:119-128.
8. Iqbal J, Pompolo S, Considine RV, Clarke IJ. Localization of leptin receptor-like immunoreactivity in the corticotropes, somatotropes, and gonadotropes in the ovine anterior pituitary. *Endocrinology* 2000; 141:1515-1520.
9. Yu WH, Kimura M, Walczewska A, Karanth S, McCann SM. Role of leptin in hypothalamic-pituitary function. *Proc Natl Acad Sci USA* 1997; 94:1023-1028.
10. Woller M, Tessmer S, Neff D, Nquema AA, Roo BV, Waechter-Brulla D. Leptin stimulates gonadotropin releasing hormone release from cultured intact hemihypothalami and enzymatically dispersed neurons. *Exp Biol Med* 2001; 226:591-596.
11. Nagatani S, Zeng Y, Keisler DH, Foster DL, Jaffe CA. Leptin regulates pulsatile luteinizing hormone and growth hormone secretion in the sheep. *Endocrinology* 2000; 141:3965-3975.
12. Amstalden M, Garcia MR, Williams SW, Stanko RL, Nizielski SE, Morrison CD, Keisler DH, Williams GL. Leptin gene expression, circulating leptin, and luteinizing hormone pulsatility are acutely responsive to short-term fasting in prepubertal heifers: relationships to circulating insulin and insulin-like growth factor I. *Biol Reprod* 2000; 63:127-133.

13. Maciel MN, Zieba DA, Amstalden M, Keisler DH, Neves J, Williams GL. Recombinant leptin prevents fasting-mediated reductions in pulsatile LH release and stimulates GH secretion in peripubertal heifers. *Proc MW Section, Am Soc Anim Sci* 2003; 66 (**Abst 263**).
14. McCann JP, Hansel W. Relationship between insulin and glucose metabolism and pituitary-ovarian functions in fasted heifers. *Biol Reprod* 1986; 34:630-641.
15. Kadokawa H, Yamada Y. Enhancing effect of acute fasting on ethanol suppression of pulsatile luteinizing hormone release via an estrogen-dependent mechanism in Holstein heifers. *Theriogenology* 1999; 51:673-680.
16. Amstalden M, Garcia MR, Stanko RL, Nizielski SE, Morrison CD, Keisler DH, Williams GL. Central infusion of recombinant ovine leptin normalizes plasma insulin and stimulates a novel hypersecretion of luteinizing hormone after short-term fasting in mature beef cows. *Biol Reprod* 2002; 66:1555-1561.
17. Zieba DA, Amstalden M, Maciel MN, Keisler DH, Raver N, Gertler A, Williams GL. Divergent effects of leptin on luteinizing hormone and insulin secretion are dose dependent. *Exp Biol Med* 2003; 228:325-330.
18. Gazal OS, Leshin LS, Stanko RL, Thomas MG, Keisler DH, Anderson LL, Williams GL. Gonadotropin-releasing hormone secretion into third-ventricle cerebrospinal fluid of cattle: correspondence with the tonic and surge release of luteinizing hormone and its tonic inhibition by suckling and neuropeptide Y. *Biol Reprod* 1998; 59:676-683.
19. Day ML, Imakawa K, Garcia-Winder M, Kittok RJ, Schanbacher BD, Kinder JE. Influence of prepubertal ovariectomy and estradiol replacement therapy on secretion of luteinizing hormone before and after pubertal age in heifers. *Domest Anim Endocrinol* 1986; 3:17-25.

20. National Research Council. Nutrient requirements of beef cattle 7th ed. Washington, D.C.: National Academy Press; 1996.
21. Zieba DA, Amstalden M, Morton S, Gallino JL, Edwards JF, Harms PG, Williams GL. Effects of leptin on basal and GHRH-stimulated GH secretion from the bovine adenohypophysis are dependent upon nutritional status. *J Endocrinol* 2003; **177:000-000 (in press)**.
22. Waters WW, Chen PL, McArthur NH, Moreno PA, Harms PG. Calcium/Calmodulin-dependent protein kinase II involvement in releasing of gonadotropin-releasing hormone. *Neuroendocrinology* 1998; 67:145-152.
23. McVey WR Jr, Williams GL. Mechanical masking of neurosensory pathways at the calf-teat interface: endocrine, reproductive, and lactational features of the suckled anestrous cow. *Theriogenology* 1991; 35:931-941.
24. Talavera F, Park CS, Williams GL. Relationships among dietary lipid intake, serum cholesterol, and ovarian function in Holstein heifers. *J Anim Sci* 1985; 60:1045-1051.
25. Kushler RH, Brown MB. A model for the identification of hormone pulses. *Stat Med* 1991; 10:329-340.
26. Cagampang FRA, Maeda KI, Yokoyama A, Ota K. Effect of food deprivation on the pulsatile LH release in the cycling and ovariectomized female rat. *Horm Metab Res* 1990; 22:269-272.
27. Samuels MH, Kramer P. Differential effects of short-term fasting on pulsatile thyrotropin, gonadotropin, and alpha-subunit secretion in healthy men – a clinical research center study. *J Clin Endocrinol Metab* 1996; 81:32-36.

28. Cameron JL, Nosbich C. Suppression of pulsatile luteinizing hormone and testosterone secretion during short-term food restriction in the adult male rhesus monkey (*Macaca mulatta*). *Endocrinology* 1991; 128:1532-1540.
29. F'Anson H, Manning JM, Herbosa CG, Pelt J, Friedman CR, Wood RI, Bucholtz DC, Foster DL. Central inhibition of gonadotropin-releasing hormone secretion in the growth-restricted hypogonadotropic female sheep. *Endocrinology* 2000; 141:520-527.
30. Kamohara S, Burcelin R, Halaas JL, Friedman JM, Charron MJ. Acute stimulation of glucose metabolism in mice by leptin treatment. *Nature* 1997; 389:374-377.
31. McArdle CA, Franklin J, Green L, Hislop JN. Signaling, cycling and desensitization of gonadotrophin-releasing hormone receptors. *J Endocrinol* 2002; 173:1-11.
32. Bjorbaek C, Uotani S, da Silva B, Flier JS. Divergent signaling capacities of the long and short isoforms of the leptin receptor. *J Biol Chem* 1997; 272:32686-32695.
33. Niswender KD, Morton GJ, Stearns WH, Rhodes CJ, Myers Jr. MG, Schwartz MW. Key enzyme in leptin-induced anorexia. *Nature* 2001; 413:794-795.
34. Takekoshi K, Ishii K, Kawakami Y, Isobe K, Nanmoku T, Nakai T. Ca²⁺ mobilization, tyrosine hydroxylase activity, and signaling mechanisms in cultured porcine adrenal medullary chromaffin cells: effects of leptin. *Endocrinology* 2001; 142:290-298.
35. Zorec R. Calcium signaling and secretion in pituitary cells. *Trends Endocrinol Metab* 1996; 7:384-388.
36. Haisenleder DJ, Dalkin AC, Marshall JC. Regulation of gonadotropin gene expression. In: Knobil E, Neill JD (eds.), *The Physiology of Reproduction*, vol. 1, 2nd ed. New York: Raven Press Ltd.; 1994:1793-1813.

37. Skinner DC, Caraty A, Malpaux B, Evans NP. Simultaneous measurement of gonadotropin-releasing hormone in the third ventricular cerebrospinal fluid and hypophyseal portal blood of the ewe. *Endocrinology* 1997; 138:4699-4704.
38. Barb CR, Chang WJ, Leshin LS, Rampacek GB, Kraeling RR. Opioid modulation of gonadotropin releasing hormone release from the hypothalamic preoptic area in the pig. *Domest Anim Endocrinol* 1994; 11:375-382.
39. Convey EM, Kesner JS, Padmanabhan V, Carruthers TD, Beck TW. Luteinizing hormone releasing hormone-induced release of luteinizing hormone from pituitary explants of cows killed before or after oestradiol treatment. *J Endocrinol* 1981; 88:17-25.
40. Fowler PA, Cunningham P, Fraser M, MacGregor F, Byrne B, Pappas A, Messinis IE, Templeton A. Circulating gonadotrophin surge-attenuating factor from superovulated women suppresses in vitro gonadotrophin-releasing hormone self-priming. *J Endocrinol* 1994; 143:45-54.
41. Henry BA, Goding JW, Alexander WS, Tilbrook AJ, Canny BJ, Dunshea F, Rao A, Mansell A, Clarke IJ. Central administration of leptin to ovariectomized ewes inhibits food intake without affecting the secretion of hormones from the pituitary gland: evidence for a dissociation of effects on appetite and neuroendocrine function. *Endocrinology* 1999; 140:1175-1182.
42. Morrison CD, Daniel JA, Holmberg BJ, Djiane J, Raver N, Gertler A, Keisler DH. Central infusion of leptin into well-fed and undernourished ewe lambs: effects on feed intake and serum concentrations of growth hormone and luteinizing hormone. *J Endocrinol* 2001; 168:317-324.

43. Morrison CD, Wood R, McFadin EL, Whitley NC, Keisler DH. Effect of intravenous infusion of recombinant ovine leptin on feed intake and serum concentrations of GH, LH, insulin, IGF-I, cortisol, and thyroxine in growing prepubertal ewe lambs. *Domest Anim Endocrinol* 2002; 22:103-112.
44. Caro JF, Kolaczynski JW, Nyce MR, Ohannesian JP, Opentanova I, Goldman WH, Lynn RB, Zhang PL, Sinha MK, Considine RV. Decreased cerebrospinal-fluid/serum leptin ration in obesity: a possible mechanism for leptin resistance. *Lancet* 1996; 348:159-161.
45. Considine RV, Sinha MK, Heiman ML, Kriauciunas A, Stephens TW, Nyce MR, Ohannesian JP, Marco CC, McKee LJ, Bauer TL et al. Serum immunoreactive-leptin concentrations in normal-weight and obese humans. *New Engl J Med* 1996; 334:292-295.
46. Halaas JL, Boozer C, Blair-West J, Fidahusein N, Denton DA, Friedman JM. Physiological response to long-term peripheral and central leptin infusion in lean and obese mice. *Proc Natl Acad Sci* 1997; 94:8878-8883.
47. Widdowson PS, Upton R, Buckingham R, Arch J, Williams G. Inhibition of food response to intracerebroventricular injection of leptin is attenuated in rats with diet-induced obesity. *Diabetes* 1997; 46:1782-1785.
48. Bjorbaek C, El-Haschimi K, Frantz JD, Flier JS. The role of SOCS-3 in leptin signaling and leptin resistance. *J Biol Chem* 1999; 274:30059-30065.
49. Emilsson V, Arch JR, de Groot RP, Lister CA, Cawthorne MA. Leptin treatment increases suppressors of cytokine signaling in central and peripheral tissues. *FEBS Lett* 1999; 455:170-174.

FIGURE LEGENDS

FIG. 1. Timeline for experimental procedures for AP (A) and HYP (B) explant incubations. Adenohypophyseal explants were perfused with KRB for 6.5 h. The first 2.5 h of perfusion served as an equilibration period (Period I). Beginning at 2.5 h, explants were perfused with KRB containing 0, 5, 10, 50, or 100 ng/ml leptin for 2 h (Period II – Basal Release). At 4.5 h, explants were challenged with 50 ng of GnRH and perfused for an additional 2 h (Period III – GnRH Stimulated). Hypothalamic-infundibular explants were equilibrated in 0.7 ml KRB for 2.5 h. Beginning at 2.5 h, explants were incubated in KRB alone or containing 1000 ng leptin for 1 h. Explants were treated at 4 h of culture with KRB containing 60 mM K^+ to induce release of GnRH as a test of viability.

FIG. 2. Mean concentrations of LH in media collected from AP explants obtained from normal-fed (top) and fasted (bottom) cows and perfused (2.5 to 4.5 h) with KRB alone (control) or KRB containing 5, 10, 50, and 100 ng/ml leptin (combined). No effect ($P > 0.1$) of leptin was observed on the release of LH (Basal Release) from AP explants from normal-fed cows (top inset). However, basal release of LH was elevated ($* P < 0.02$) in leptin-treated AP explants from fasted cows (bottom inset). Arrows indicate the beginning of control (KRB alone) or leptin treatment.

FIG. 3. Mean concentrations of LH in media after GnRH stimulation of AP explants of normal-fed and fasted cows perfused in the presence of either KRB alone (control) or KRB containing 5, 10, 50, and 100 ng/ml leptin. All doses of leptin increased ($P < 0.001$) GnRH-stimulated release of LH in AP explants collected from normal-fed (top panel), but not fasted (bottom

panel) cows. Adenohypophyseal explants from fasted cows perfused with 5 ng/ml leptin released less ($P < 0.001$) LH in response to GnRH than control-treated explants. Note the elevated baseline for leptin-treated compared to control-treated AP explants from fasted cows at the time of GnRH stimulation (4.5 h).

FIG. 4. Fold change in mean concentrations of GnRH in media collected from HYP explants of normal-fed and fasted cows. Hypothalamic-infundibular explants were incubated in 0.7 ml of KRB either alone or containing 1000 ng leptin at 2.5 and 3 h. Explants were incubated with KRB containing 60 mM K^+ at 4 h of incubation to test viability of tissue. No effects of diet or leptin were observed ($P > 0.1$).

TABLE 1. Means (\pm SEM) concentrations, and frequency and amplitude of LH pulses in normal-fed and fasted cows.

Parameter	Normal Fed		Fasted	
	Day 0	Day 3	Day 0	Day 3
Concentration (ng/ml)	5.2 \pm 0.1	5.9 \pm 0.1	4.2 \pm 0.1	5.3 \pm 0.2
Frequency (pulses/6 h)	8.3 \pm 1.2	10 \pm 1.0	5.0 \pm 1.4	7.0 \pm 1.6
Amplitude (ng/ml)	1.6 \pm 0.6	2.0 \pm 0.5	1.7 \pm 0.8	1.3 \pm 0.2

No dietary effect was observed.




FIG. 1




FIG. 2




FIG. 3




FIG. 4